

## Week 10: Lecture 1

**Non-Reacting Mixtures****Definitions**

- the total mass of a mixture

$$m = m_1 + m_2 + \cdots + m_j = \sum_{i=1}^j m_i$$

- the relative amount of components specified in terms of mass fractions

$$Y_i = \frac{m_i}{m} \Rightarrow \sum_{i=1}^j Y_i = 1$$

- the total number of moles of a mixture

$$n = n_1 + n_2 + \cdots + n_j = \sum_{i=1}^j n_i$$

- the relative amount of components specified in terms of mole fractions

$$X_i = \frac{n_i}{n} \Rightarrow \sum_{i=1}^j X_i = 1$$

- $m_i$  and  $n_i$  are related by the molecular weight  $\tilde{M}_i$

$$m_i = n_i \cdot \tilde{M}_i$$

therefore

$$m = \sum_{i=1}^j n_i \cdot \tilde{M}_i$$

**Week 10: Lecture 1**

- the mixture molecular weight can be calculated as a mole-fraction average of the component molecular weights

$$\tilde{M} = \frac{m}{n} = \frac{\sum_{i=1}^j n_i \tilde{M}_i}{n} = \sum_{i=1}^j X_i \tilde{M}_i$$

- $X_i$  and  $Y_i$  are also related by the molecular weight

$$\frac{Y_i}{X_i} = \frac{m_i/m}{n_i/n} = \left( \frac{m_i}{n_i} \right) \left( \frac{m}{n} \right) = \left( \tilde{M}_i \right) \left( \frac{1}{\tilde{M}} \right)$$

therefore

$$Y_i = X_i \left[ \frac{\tilde{M}_i}{\sum_{i=1}^j X_i \tilde{M}_i} \right]$$